



**OcTTain™ XL**

## QUICK FACTS

An easy-to-use new level of broadleaf weed control, kochia and wild buckwheat in barley, wheat and durum.

<b>ACTIVE INGREDIENT:</b>	2,4-D Ester and Fluroxypyr
<b>GROUP:</b>	4
<b>PACKAGING:</b>	2 x 9 L
<b>RAINFAST:</b>	1 hour
<b>WATER VOLUME:</b>	Ground application: 5 – 10 gal/acre Air application: 3 – 5 gal/acre
<b>STORAGE:</b>	Avoid freezing. If frozen, bring to room temperature and agitate.
<b>REGISTERED CROPS:</b>	Spring Wheat, Winter Wheat, Durum Wheat, Barley, Smooth Bromegrass*, Meadow Bromegrass*, Timothy*, Crested Wheatgrass*, Intermediate Wheatgrass*, Creeping Red Fescue*, Tall Fescue* *Seedling and established for seed production only
<b>APPLICATION RATE:</b>	0.45 L/acre (40 acres/case)
<b>TANK MIX OPTIONS:</b>	Simplicity™, Fenoxaprop (Bengal® WB, Cordon®, Vigil WB), Tralkoxydim (Liquid Achieve™, Bison®, Nufarm Tralkoxydim Liquid), Clodinafop (Ladder® 240EC, Slam'R®, Signal®), 2,4-D Ester *Use any surfactant required for tank-mix partner





# TANK MIX ORDER

Add grassy tank-mix partner first\*\* followed by Octtain XL followed by any adjuvant / surfactant if required.

\*\* Fenoxaprop and Traxos tankmix partners should go in the tank after Octtain XL

# KEY WEEDS CONTROLLED\*

- Cleavers
- Flixweed
- Hemp-nettle
- Kochia (group 2 and 9 resistant)
- Lamb's Quarters
- Round-leaved Mallow
- Shepherd's-purse
- Stinkweed
- Stork's-bill
- Volunteer Canola (all biotypes)
- Volunteer Flax
- Wild Buckwheat
- Wild Mustard

\*Refer to product label for detailed weed staging information.

# PRO TIPS

- Spring crops apply from 4 leaf to emergence of flag leaf
- Winter wheat apply from 3 tiller stage to just prior to flag leaf stage
- For maximum performance apply between 12 – 24 °C
- 60 day pre-harvest interval
- Follow grazing restrictions if crop is intended for feed use

## ALWAYS READ AND FOLLOW LABEL DIRECTIONS

\*Registered trademarks of Federated Co-operatives Limited