

## **QUICK FACTS**

A selective post-emergent herbicide for the control of annual and perennial grasses in canola, flax and pulses, and is an excellent option for controlling foxtail barley.

**ACTIVE INGREDIENT:** Quizalofop P-ethyl

GROUP:

**PACKAGING:** 1 x 8 L Contender and 1 x 8 L Adjuvant

RAINFAST: 1 hour

**WATER VOLUME:** Ground application: 10 gal/acre (use higher rate to improve coverage)

Aerial application: 3 – 5 gal/acre

**STORAGE:** Do not freeze

**REGISTERED CROPS:** Alfalfa (for seed), Camelina, Canola, Canola-Quality Brassica Juncea, Chickpeas,

Creeping Red Fescue (for seed), Dry Beans, Established Alsike Clover (for seed), Established Red Clover (for seed), Faba Beans, Flax, Lentils, Mustard, Narrow Leaf Lupin, Peas, Seedling Legume (for seed), Soybeans, Sugar Beets, Sunflowers

0.15 – 0.30 L/acre (26 – 53 acres/case)

**TANK MIX OPTIONS:** Ally<sup>®</sup>, Basagran<sup>®</sup>, Basagran<sup>®</sup> Forté, Express<sup>®</sup> SG, Muster<sup>®</sup>, Pinnacle<sup>®</sup>

\*Apply tank-mix partner to registered crops only



**APPLICATION RATE:** 



### TANK MIX ORDER

Add tank mix partner first (except Basagran Forté) followed by Contender herbicide, then adjuvant (add Basagran Forté after contender but before adjuvant).

# **KEY WEEDS CONTROLLED\***

- Barnyard Grass
- · Downy Brome
- Foxtail Barley
- · Green Foxtail

- Japanese Brome
- Oats
- Quackgrass
- Volunteer Barley

- Volunteer Corn
- Volunteer Wheat
- Wild Oats
- Yellow Foxtail

#### **PRO TIPS**

- Use 50 mesh screens or larger. A spray pressure of 210 275 kPa (30 40 psi) is recommended. Avoid overlapping. Shut off spray booms while starting, turning, slowing or stopping to prevent over application.
- Use spray preparation of IPCO Contender Herbicide within 24 hours or product degradation may occur. If spray preparation is left without agitation, thoroughly agitate before using.
- · Do not apply when environmental conditions such as wind speed, temperature, relative humidity, etc. are favourable for drift to occur.
- For the majority of weeds, apply at the 2 3 leaf stage for optimum control.

#### ALWAYS READ AND FOLLOW LABEL DIRECTIONS

®Registered trademarks of Federated Co-operatives Limited

<sup>\*</sup>Ensure appropriate rate is selected based on weed type and weed stage.