



# PATRON 240 EC

Broad-spectrum grassy weed control in canola, flax and pulses.



## **QUICK FACTS:**

Active Ingredient Clethodim (240 g/L)

Group

**Packaging** One case (3 L jug of Patron + 9 L

of adjuvant) = 20-60 acres

Rainfast 1 hour

Water Volume Minimum of 22 L/acre when

applied alone (under ideal conditions)

**Re-cropping** None

Restrictions

Storage

May be stored at any temperature.

### **REGISTERED CROPS**

Alfalfa, seedlings Canola Chickpeas Coriander Dry beans Fenugreek Field Peas
Flax Lentils Mustard Potatoes Soybeans Sunflowers

### **MIXING INSTRUCTIONS:**

- 1. Fill clean spray tank 1/2 full with water; start agitation
- 2. Add the correct amount of Patron 240EC herbicide and continue to agitate
- 3. Add the correct amount of Patron adjuvant
- 4. Fill the spray tank with the remaining amount of water
- 5. Continue to agitate or run the bypass system
- 6. If there is a break in the spraying operation, agitate the mixture thoroughly before spraying
- 7. Do not allow the mixture to sit overnight
- 8. If tank mixing, always follow label directions for each tank-mix partner

**Quick Tip:** Clethodim activity may be reduced by levels of bicarbonates over 500 ppm in the water. The addition of ammonium sulphate (AMS) at 1% v/v can help restore control.

SHAKE WELL BEFORE USING. Always read and follow label directions.

### WEEDS CONTROLLED\*

Weed		Application Stage	Recommended Rate	
			mL per acre	acres per case
Foxtail (green, yellow) Wild oats	Volunteer cereals (wheat, barley, oats)	2–4 leaf stage	50	60
Barnyard grass Proso millet	Volunteer corn Volunteer canary grass	2–6 leaf stage	50	60
Wild oats Volunteer cereals (wheat, barley, oats) Foxtail (green, yellow) Persian darnel Crabgrass (smooth and large)	Proso millet Witchgrass Fall panicum Barnyard grass Volunteer corn Volunteer canary grass	2–6 leaf stage	80	40
Quackgrass control		2–6 leaf stage	150	20

<sup>\*</sup>PATRON 240 EC applied at a rate of 60 acres per case should only be applied under the following conditions:

- $\bullet \operatorname{\mathsf{Good}} \operatorname{\mathsf{crop}} \operatorname{\mathsf{stand}}$
- When not tank-mixed with other pesticides
- Light to moderate weed infestation
- Absence of stress, good growing conditions
- Early application (as in above table, prior to tillering)
- Do not apply to volunteer winter cereals
- · Adequate moisture and fertility

If any one of the above conditions is not present at the time of application, use the 40 acre per case rate of PATRON 240 EC.

PATRON™ ™Trademark of Federated Co-operatives Limited, Saskatoon S7K 3M9 CO-OP® ©CO-OP and design trademark are registered trademarks of TMC Distributing Ltd., Saskatoon S7K 3M9.

