



2,4-D

ESTER 700

QUICK FACTS

A post-emergent broadleaf herbicide in spring wheat, barley, rye and winter wheat.

ACTIVE INGREDIENT:	2,4-D
GROUP:	4
PACKAGING:	2 x 10 L
RAINFAST:	2 hours
WATER VOLUME:	Ground application: 5 – 10 gal/acre Aerial application: 3 gal/acre
STORAGE:	May be frozen
REGISTERED CROPS:	Barley, wheat (spring, winter), established pasture grass, grass crops for forage and seed, rye (spring, fall), sod and turf
APPLICATION RATE:	Annual Crops: 192 – 445 ml/acre Forage Grass and Pasture: 607 – 1012 ml/acre *Ensure appropriate rate is selected based on crop type
TANK MIX OPTIONS:	Express® SG, Refine® SG, Ally® *Other products may have other 2,4-D tank mix options on their label





TANK MIX ORDER

When used in a tank mix, consult product label(s) and use W.A.L.E.S guidelines.

KEY WEEDS CONTROLLED*

- Annual sow thistle
- Biennial wormwood
- Bluebur (up to 4-leaf stage)
- Canada thistle
- Chickweed
- Common ragweed
- Curled dock
- Dandelion
- Field bindweed
- Field horsetail
- Flixweed
- Goat's beard
- Hawkweed
- Hemp-nettle
- Knotweed
- Lady's thumb
- Lamb's quarters
- Narrow-leaved hawk's beard
- Peppergrass
- Perennial sow-thistle
- Pineapple weed
- Plantain
- Prickly lettuce
- Prostrate pigweed
- Purslane
- Redroot pigweed
- Russian knapweed
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Sweet clover (seedling)
- Volunteer canola
- Volunteer sunflower
- Wild buckwheat
- Wild Mustards
- Wild radish
- Wild (prairie) sunflower

Please see label for additional pasture weeds controlled.

*Ensure appropriate rate is selected based on weed type and weed stage.

PRO TIPS

- For annual cereals, apply from the 4-leaf stage up to just before flag leaf
- Allow 30 days prior to harvesting forage
- May also be applied for vegetation management purposes on non-crop land

ALWAYS READ AND FOLLOW LABEL DIRECTIONS

*Registered trademarks of Federated Co-operatives Limited